

14-05

## STATEMENT OF POLICY

### Vector-Borne Disease

#### Policy

The National Association of County and City Health Officials (NACCHO) strongly urges the federal government to provide sufficient funds to maintain, strengthen, and expand the surveillance, disease prevention, and research capacities necessary to identify, track, and address existing and emerging vector-borne diseases (VBDs).

NACCHO supports local public health activities to prevent, monitor, and control such diseases, including the following:

- Working collaboratively with local, state, and national partners from multiple disciplines.
- Training local health department staff to investigate VBD cases in humans and animals, collect vector samples, and perform abatement.
- Expanding laboratory capacity to develop and maintain testing capacity for established and emerging vector-borne pathogens in human, animal, and vector samples.
- Improving data collection systems for identifying and tracking VBD cases and distribution in humans and animals.
- Increasing the effective use of human and animal surveillance data to target prevention and control efforts.
- Enhancing data-sharing systems to facilitate effective communication and monitoring.
- Participating in longitudinal monitoring programs to study changes in vector distribution and density over time.
- Developing prevention and control plans to reduce the impact of established and emerging vector-borne diseases on local communities, including vulnerable populations, and to address environmental sources of emerging infectious diseases through a “One Health” approach.<sup>1</sup>
- Educating the public through social media, local media, and educational materials.
- Educating healthcare providers to improve VBD diagnosis and reporting.

Effectively addressing VBDs will also require consideration of related topics covered in NACCHO’s policy statements [Climate Change](#) and [Mosquito Control](#).

#### Justification

Vector-borne diseases are diseases transmitted by insects to humans and domestic animals. These diseases can lead to significant morbidity, including long-term sequelae and mortality in the United States. Surveillance for these vectors and human/animal disease is necessary for prevention and control. There is also a critical need to maintain and strengthen local public health systems’ capacity to prevent cases and contain outbreaks of VBDs through effective



abatement efforts, public and provider education, and preparedness planning. Reductions in vector surveillance and prevention capacity compromise local and national ability to detect changes in vector activities, as well as mitigate the impact of established and emerging vector-borne diseases on local communities.

The need for a robust vector surveillance infrastructure is critical in reducing the impact of a variety of VBDs. Lyme disease is endemic in North America and since standardized reporting began in the U.S., case counts have increased steadily. The U.S. is seeing an expansion in the geographic range of tick-borne Powassan virus. In 2012, the nation experienced one of the most intense outbreaks of West Nile Virus (WNV), resulting in 286 deaths.<sup>2</sup> Dengue has also established itself in the continental U.S., with transmission documented in Florida, Texas, and New York.<sup>3</sup> The U.S. periodically experiences outbreaks of mosquito-borne encephalitis viruses including Eastern, LaCrosse, St. Louis, and California Group encephalitis. These and other emerging vector-borne diseases pose a significant threat to public health.<sup>4</sup>

In addition, the appearance of the Zika virus in the Americas in 2014 presented an unprecedented challenge for public health systems. The rapid spread of the virus in mosquitoes and its newly-detected association with severe birth defects illustrate how vector-borne diseases may move rapidly and may change in their clinical behavior. Similar to trends seen among ticks, mosquitoes are expanding their range. After its introduction into the U.S. in the mid-1980s, *Aedes albopictus* (Asian Tiger mosquito), which can transmit several viruses including Yellow Fever, Dengue, Chikungunya, and Zika viruses, has been expanding its range throughout the southeastern U.S. into the northeast and western parts of the country.<sup>5</sup>

Prior to the introduction of WNV in the U.S. in 1999, there was no federal support for state or local VBD surveillance infrastructure.<sup>6</sup> In response to the introduction and subsequent spread of WNV, approximately \$24 million federal dollars was awarded to states and the six largest cities/counties that year. A survey in 2004 showed this funding had created a robust and well-integrated national surveillance infrastructure for arboviruses, a common type of VBD. The surveillance system that was developed, ArboNET, depends on state and local health departments to submit human case information, mosquito testing data, and laboratory testing data. However, from 2004 to 2012, annual federal arbovirus surveillance funding declined to \$9.3 million, a reduction that negatively affected national surveillance capacity. A survey of the arbovirus surveillance infrastructure—conducted in 2013 and compared to the 2004 survey—showed substantial erosion of public health capacity to detect disease and mosquito infection and conduct essential laboratory testing.<sup>2,6,7</sup> The survey revealed that active surveillance for human cases declined by 22% and mosquito surveillance declined by 9%. Further, NACCHO's 2017 assessment, [\*Mosquito Capabilities in the U.S.\*](#), revealed that about 46% of vector control programs are not conducting routine surveillance for mosquitoes. Overall, the assessment identified 84% of vector control programs as needing improvement.<sup>7,9</sup> Reductions in mosquito surveillance affect health departments' ability to understand the mosquito populations in their area and make the best decisions regarding control measures.

Each of these reductions affects local and national ability to measure the risk of VBDs. Strong surveillance systems, technical expertise, laboratory capacity, and prevention programs are also needed to prevent and control the spread of established and emerging VBDs. Given that current

vector and vector-borne disease surveillance and research efforts remain underfunded, it is vital that additional funds be allocated to maintain, strengthen, and expand surveillance and research capacities. Past reductions in public health funding compromise the capacity of local health departments to conduct surveillance and control of mosquitoes, other vectors, and associated diseases.<sup>6,7,10</sup> The public health infrastructure must be strengthened if we hope to sustain surveillance; detect emerging diseases; and prevent disease spread through effective vector control and behavioral change strategies to protect communities against these serious and potentially devastating diseases.

## **References**

1. CDC. (2013). *One Health*. Retrieved February 15, 2018, from <http://www.cdc.gov/onehealth/>
2. CDC. (2014). National capacity for surveillance, prevention, and control of West Nile Virus and other arbovirus infections – United States, 2004 and 2012. *Morbidity and Mortality Weekly Report*; 63(13):281-284.
3. CDC. (2010). Locally-acquired dengue – Key West, Florida, 2009-2010. *MMWR*; 59:577-581.
4. Fauci AS, Morens DM. Zika Virus in the Americas – Yet Another Arbovirus Threat. *N Engl J Med* 2016; 374:601-604. February 18, 2016. DOI: 10.1056/NEJMp1600297
5. Ruiz-Moreno, D., Vargas, I.S., Olson, K.E., & Harrington, L.C. (2012). Modeling dynamic introduction of Chikungunya virus in the United States. *PLoS Negl Trop Dis*;6:e1918.
6. CDC. (2006). Assessing capacity for the surveillance, prevention and control of West Nile Virus infection – United States, 1999-2004. *Morbidity and Mortality Weekly Report*; 55:150-153.
7. Hadler, J. L., Patel, D., Nasci, R. S., Petersen, L. R., Hughes, J. M., Bradley, K., Engel, J. (2015). Assessment of Arbovirus Surveillance 13 Years after Introduction of West Nile Virus, United States. *Emerging Infectious Diseases*, 21(7), 1159-1166. Retrieved November 2, 2017 from <https://dx.doi.org/10.3201/eid2107.140858>.
8. National Association of County and City Health Officials (NACCHO). (2017). *Mosquito Control Capabilities in the U.S.* Washington, DC. Retrieved January 22, 2018 from <https://www.naccho.org/uploads/downloadable-resources/Mosquito-control-in-the-U.S.-Report.pdf>
9. National Association of County and City Health Officials (NACCHO). (2016). *Mosquito Surveillance and Control Assessment in Zika Virus Priority Jurisdictions*. Washington, DC. Retrieved January 22, 2018 from <https://www.naccho.org/uploads/downloadable-resources/VectorAssessmentDec2016NACCHO.pdf>
10. CDC. (2013). *West Nile Virus in the United States: Guidelines for Surveillance, Prevention and Control*. 4<sup>th</sup> revision. Fort Collins, CO: US Department of Health and Human Services. Retrieved June 11, 2014, from <http://www.cdc.gov/westnile/resources/pdfs/wnvguidelines.pdf>

## **Record of Action**

*Proposed by NACCHO Epidemiology and Infectious Disease Prevention and Control Workgroups*

*Approved by NACCHO Board of Directors July 8, 2014*

*Updated March 2018*