

Characteristics of Medications for Opioid-Addiction Treatment.

Characteristic	Methadone	Buprenorphine	Naltrexone
Brand names	Dolophine, Methadose	Subutex, Suboxone, Zubsolv	Depade, ReVia, Vivitrol
Class	Agonist (fully activates opioid receptors)	Partial agonist (activates opioid receptors but produces a diminished response even with full occupancy)	Antagonist (blocks the opioid receptors and interferes with the rewarding and analgesic effects of opioids)
Use and effects	Taken once per day orally to reduce opioid cravings and withdrawal symptoms	Taken orally or sublingually (usually once a day) to relieve opioid cravings and withdrawal symptoms	Taken orally or by injection to diminish the reinforcing effects of opioids (potentially extinguishing the association between conditioned stimuli and opioid use)
Advantages	High strength and efficacy as long as oral dosing (which slows brain uptake and reduces euphoria) is adhered to; excellent option for patients who have no response to other medications	Eligible to be prescribed by certified physicians, which eliminates the need to visit specialized treatment clinics and thus widens availability	Not addictive or sedating and does not result in physical dependence; a recently approved depot injection formulation, Vivitrol, eliminates need for daily dosing
Disadvantages	Mostly available through approved outpatient treatment programs, which patients must visit daily	Subutex has measurable abuse liability; Suboxone diminishes this risk by including naloxone, an antagonist that induces withdrawal if the drug is injected	Poor patient compliance (but Vivitrol should improve compliance); initiation requires attaining prolonged (e.g., 7-day) abstinence, during which withdrawal, relapse, and early dropout may occur