

## PrEP Providers List – Public Health – Seattle & King County

The following is a list of medical providers who have expressed a willingness to screen for, prescribe, and monitor pre-exposure prophylaxis (PrEP) for people at high risk of HIV infection.

### Medical Providers:

Provider(s)	Clinic	Phone
Dave Holt, ARNP Jessica Rongitsch, MD Vy Chu, MD	Capitol Hill Medical	206 720 9999
Karen Johnson, ARNP Katie Hester ARNP Matthew J. Logalbo, MD Katherine Mayer, MD Sonja Olson, MD	Country Doctor Community Clinic	206 299 1600
Kathy Brown, MD	Group Health Cooperative Burien, Capitol Hill, Downtown, & Northgate	206 326 3609
Byron Doepker, MD Greg Allen, MD Kim Waarvick, MD	Group Health Cooperative Capitol Hill	206 326 3530
Alison Roxby, MD Bob (Robert) Harrington, MD Connie Celum, MD, MPH H. Nina Kim, MD, MSc Jared Baeten, MD, PhD Jeanne Marrazzo, MD, MPH Mary Campbell, MD Ruanne Barnabas, MD Shireesha Dhanireddy, MD	Madison Clinic at Harborview	206 744 5100
Paul Algeo, PharmD, PA-C Peter Shalit, MD, PhD	Peter Shalit's Office	206 624 0688
Amy Treakle, MD	Polyclinic Infectious Diseases	206 860 4531
Trong Tony Trinh, MD, MPH		206 860 4447
Michael Bolton, MD, PhD		206 860 4603
Joanne Stekler, MD, MPH	PrEP Clinic at Gay City Wellness Center	206 860-6969
Emma Swingle, MD	Qliance	206 913 4700
Rob Killian, MD	Rob Killian's Office	206 568 6320
Alison Roxby, MD Ann Stapleton, MD Bob (Robert) Coombs, MD, PhD David Koelle, MD Geoff Gottlieb, MD, PhD	Virology Clinic at UWMC-Roosevelt	206 598 8750
David Aboulafia, MD	Virginia Mason Medical Center Hematology/Oncology	206 223 6193
Chia Wang, MD Chris Baliga, MD Margo Schwartz, MD Shingo Chihara, MD Uma Malhotra, MD	Virginia Mason Medical Center (Infectious Diseases)	206 341-0846
Nancy Connolly, MD, MPH	Virginia Mason Medical Center, Lynnwood	425 712 7900

Warren Dinges, MD, PhD	Warren Dinges Practice: Seattle Infectious Disease Clinic	206 682 3444
------------------------	---	--------------

**For assistance in paying for PrEP, please see the Washington State PrEP Drug Assistance Program (PrEP DAP)**

<http://www.doh.wa.gov/YouandYourFamily/IllnessandDisease/HIVAIDS/HIVCareClientServices/PrEP DAP.aspx>

**For Providers:**

The PrEP Providers list includes HIV medical providers who have expressed interest in evaluating patients for HIV pre-exposure prophylaxis (PrEP) and whom are willing to prescribe PrEP to persons at high-risk for HIV infection. Public Health maintains this list on our website and medical providers in the community may be added to or deleted from the list at any time by sending an email to [buskins@kingcounty.gov](mailto:buskins@kingcounty.gov)

Medical providers on the list should be willing to follow the published CDC PrEP guidelines. These guidelines include that:

- 1) PrEP is contraindicated in persons with unknown or positive HIV status or with an estimated creatinine clearance <60 mL/min;
- 2) PrEP should be targeted to adults at very high risk for HIV acquisition;
- 3) PrEP should be delivered as part of a comprehensive set of prevention services; and
- 4) PrEP should be accompanied by quarterly monitoring of HIV status, pregnancy status, side effects, medication adherence, and risk behaviors, as outlined in previous interim guidance.<sup>1-3</sup>

1. US Public Health Service. Preexposure prophylaxis for the prevention of HIV infection in the United States – 2014: a clinical practice guideline. Published 5/14/2014. Available at <http://www.cdc.gov/hiv/pdf/PrEPguidelines2014.pdf>

2. CDC. Interim guidance: preexposure prophylaxis for the prevention of HIV infection in men who have sex with men. MMWR 2011;60:65–8. (Updated June 14, 2013 [http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6223a2.htm?s\\_cid=mm6223a2\\_w](http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6223a2.htm?s_cid=mm6223a2_w))

3. CDC. Interim guidance for clinicians considering the use of preexposure prophylaxis for the prevention of HIV infection in heterosexually active adults. MMWR 2012;61:586–9.

Public Health **further** recommends that medical providers discuss PrEP with all HIV-uninfected men who have sex with men being treated for rectal gonorrhea or syphilis, and that providers use either a 4<sup>th</sup> generation HIV EIA or an HIV RNA test when testing persons prior to PrEP initiation or as part of PrEP monitoring. Thank you for your ongoing efforts to prevent HIV in our community.